



UC DAVIS
VETERINARY MEDICINE

California Animal Health & Food Safety Laboratory System

PO Box 1770, Davis, CA, 95617

(530) 752-8700

www.cahfs.vetmed.ucdavis.edu

CAHFS Accession #: D2201812

FINAL REPORT

Ref.#:

Coordinator: Omar Gonzales Viera, DVM, PhD,
Dipl. ACVP, Resident

E-Signed and Authorized by: Gonzales Viera,
Omar on 2/9/2022 6:37:41PM

Email To:
Cascade Valley Heritage Farm
info@cvhfarm.com

Collection Site:
Cascade Valley Heritage Farm - Priya Tuvell
Chico CA 95973
Butte County

This report supersedes all previous reports for this case

Specimens Received: 11 Clotted Blood - RTT;

Date Collected: 02/05/2022 **Date Received:** 02/08/2022

Comments: 17 rtt

Case Contacts

Submitter	Cascade Valley Heritage Farm	Chico	CA	95973
-----------	------------------------------	-------	----	-------

Specimen Details

Animal/Source	ID Type	Taxonomy	Gender	Age
1	CAHFS Internal ID	Nigerian Dwarf goat	Mixed	Adult

Laboratory Findings/Diagnosis

Negative for antibodies to Brucella, C. burnetii, C. pstb, CAE/OPP and M. avium paratuberculosis

Clinical History

Routine Annual Health Screening.

Immunology

Test Specific Comments

Caprine arthritis encephalitis / ovine progressive pneumonia antibody cELISA

- * Results in the suspect range may be caused by early-stage infection, or may be temporary cross-reaction due to recent vaccination, stress, or other illness. Re-testing the animal in 3-4 weeks is recommended. Most suspect results due to temporary cross-reaction revert to negative in this time period.

Corynebacterium pseudotuberculosis antibody - SHI

* *C. pseudotuberculosis* SHI titers less than 1:8 are not considered significant. Titers may indicate active infection or historic exposure.

In small ruminants, *C. pseudotuberculosis* titers of 1:8 occur with similar frequency in animals with and without active infection and may represent early infection or historic exposure. Titers 1:16-1:32 occur 4 times more often in animals with active infection (vs. those without infection), and titers 1:64->=1:256 occur 8.5 times more often in animals with active infection (vs. those without).

In horses, titers 1:8-1:128 are typically seen in animals with historic infection with or exposure to *C. pseudotuberculosis*, with external infection, or in the early stages of infection. Titers >=1:256 may be seen with active infection (external, internal) or exposure to the organism. In the absence of external abscesses, and with other indicators of chronic inflammation (such as high WBC, high globulin, others), titers >1:512 are associated with internal abscess formation.

M. paratuberculosis (Johne's Disease) Antibody ELISA

* In cattle, *Mycobacterium paratuberculosis* (Johne's disease) ELISA test results are interpreted as negative if less than or equal to 0.450, suspect if greater than 0.450 but less than 0.550, and positive if greater than or equal to 0.550. At the 0.550 cutoff, test sensitivity and specificity are estimated at 68% and 99%, respectively, in cattle.

In sheep and goats, *Mycobacterium paratuberculosis* (Johne's disease) ELISA test results are interpreted as negative if less than or equal to 0.100, suspect if greater than 0.100 but less than 0.300, and positive if greater than or equal to 0.300. At the 0.300 cutoff, test sensitivity and specificity are estimated at 51% and 95%, respectively, in small ruminants.

Brucella (B. melitensis) 3% Rose Bengal antibody test

Animal/Source	Specimen	Specimen Type	Results
1	(3) War Marchine	Clotted Blood - RTT	Negative
1	(8) Elysian	Clotted Blood - RTT	Negative
1	(9) Suadela	Clotted Blood - RTT	Negative
1	(10) Willow	Clotted Blood - RTT	Negative
1	(11) Starlit	Clotted Blood - RTT	Negative

Caprine arthritis encephalitis / ovine progressive pneumonia antibody cELISA

Animal/Source	Specimen	Specimen Type	Results
1	(1) Hullabaloo	Clotted Blood - RTT	8.5 Neg
1	(2) Zinqua Boom	Clotted Blood - RTT	-0.51 Neg
1	(3) War Marchine	Clotted Blood - RTT	0.42 Neg
1	(4) Aspen	Clotted Blood - RTT	-0.51 Neg
1	(5) Sunny/Sunspot	Clotted Blood - RTT	-0.98 Neg
1	(6) Juna	Clotted Blood - RTT	-0.98 Neg
1	(7) Denali	Clotted Blood - RTT	2.28 Neg
1	(8) Elysian	Clotted Blood - RTT	1.72 Neg
1	(9) Suadela	Clotted Blood - RTT	-3.11 Neg
1	(10) Willow	Clotted Blood - RTT	2.37 Neg
1	(11) Starlit	Clotted Blood - RTT	-1.63 Neg

Corynebacterium pseudotuberculosis antibody - SHI

Animal/Source	Specimen	Specimen Type	Results
1	(1) Hullabaloo	Clotted Blood - RTT	Neg @1:8
1	(2) Zinqua Boom	Clotted Blood - RTT	Neg @1:8
1	(3) War Marchine	Clotted Blood - RTT	Neg @1:8
1	(4) Aspen	Clotted Blood - RTT	Neg @1:8
1	(5) Sunny/Sunspot	Clotted Blood - RTT	Neg @1:8
1	(6) Juna	Clotted Blood - RTT	Neg @1:8

1	(7) Denali	Clotted Blood - RTT	Neg @1:8
1	(8) Elysian	Clotted Blood - RTT	Neg @1:8
1	(9) Suadela	Clotted Blood - RTT	Neg @1:8
1	(10) Willow	Clotted Blood - RTT	Neg @1:8
1	(11) Starlit	Clotted Blood - RTT	Neg @1:8

Coxiella burnetii (Q Fever) antibody ELISA

Animal/Source	Specimen	Specimen Type	Results
1	(3) War Marchine	Clotted Blood - RTT	Negative
1	(8) Elysian	Clotted Blood - RTT	Negative
1	(9) Suadela	Clotted Blood - RTT	Negative
1	(10) Willow	Clotted Blood - RTT	Negative
1	(11) Starlit	Clotted Blood - RTT	Negative

M. paratuberculosis (Johne Disease) Antibody ELISA

Animal/Source	Specimen	Specimen Type	Results
1	(1) Hullabaloo	Clotted Blood - RTT	0.003
1	(2) Zinquá Boom	Clotted Blood - RTT	0.00
1	(3) War Marchine	Clotted Blood - RTT	0.003
1	(4) Aspen	Clotted Blood - RTT	0.004
1	(5) Sunny/Sunspot	Clotted Blood - RTT	0.005
1	(6) Juna	Clotted Blood - RTT	0.001
1	(7) Denali	Clotted Blood - RTT	0.00
1	(8) Elysian	Clotted Blood - RTT	0.00
1	(9) Suadela	Clotted Blood - RTT	0.00
1	(10) Willow	Clotted Blood - RTT	0.003
1	(11) Starlit	Clotted Blood - RTT	0.00

We VALUE your opinion of our services!

To help us improve, please take a moment to tell us your impression of the services you received.
Please click the link below to be directed to a short survey

[CAHFS Client Feedback Survey](#)